

Some exact solutions for the Caudrey-Dodd-Gibbon equation

Alvaro Salas*

Abstract

In this paper we show some exact solutions for the Caudrey-Dodd-Gibbon equation (CDG equation). These solutions are obtained via $\text{\textcircled{R}}Mathematica$ 6.0 by the projective Riccati equation method.

Key words and phrases: Nonlinear differential equation, nonlinear partial differential equation, fifth order evolution equation, fifth order KdV equation, traveling wave solution, projective Riccati equation method, partial differential equation (PDE), KdV equation, Caudrey-Dodd-Gibbon equation, $\text{\textcircled{R}}Mathematica$.

1 Introduction

From 70's, a vast variety of the simple and direct methods to find analytic solutions of nonlinear differential equations and evolution equations have been developed. Recently, the projective Riccati equation method has drawn lots of interests in seeking the solitary wave solution and other kinds of solutions.

The general form of the fifth-order KdV equation is written as

$$u_t + \omega u_{xxxxx} + \alpha uu_{xxx} + \beta u_x u_{xx} + \gamma u^2 u_x = 0 \quad (1.1)$$

*Department of Mathematics, Universidad de Caldas, Department of Mathematics, Universidad Nacional de Colombia, Manizales, Colombia. *email* : asalash2002@yahoo.com

where α , β , γ and ω are arbitrary real parameters. In this paper we present exact solutions for a particular case of this equation obtained by setting $\alpha = 30$, $\beta = 30$, $\gamma = 180$ and $\omega = 1$. Thus, the equation to be studied is

$$u_t + u_{xxxxx} + 30uu_{xxx} + 30u_xu_{xx} + 180u^2u_x = 0 \quad (1.2)$$

Equation (1.2) is known as Caudrey-Dodd-Gibbon equation.

2 The projective Riccati equation method

We search exact solutions of equation (1.2) in the form

$$\begin{cases} u(x, t) = v(\xi) \\ \xi = x + \lambda t, \end{cases} \quad (2.3)$$

As a result we have that the equation (1.2) is reduced to the nonlinear ordinary differential equation (ODE)

$$180V'(\xi)V(\xi)^2 + 30V^{(3)}(\xi)V(\xi) + \lambda V'(\xi) + 30V'(\xi)V''(\xi) + V^{(5)}(\xi) = 0 \quad (2.4)$$

To obtain exact solution for the equation (2.4), we use the projective Riccati equation method [4][7] which may be described in the following three steps:

Step 1. We consider solutions of (2.4) in the form

$$v(\xi) = a_0 + \sum_{j=1}^m \sigma(\xi)^{j-1} (a_j \sigma(\xi) + b_j \tau(\xi)), \quad (2.5)$$

where $\sigma(\xi)$, $\tau(\xi)$ satisfy the system

$$\begin{cases} \sigma'(\xi) = e\sigma(\xi)\tau(\xi) \\ \tau'(\xi) = e\tau^2(\xi) - \mu\sigma(\xi) + r. \end{cases} \quad (2.6)$$

It may be proved that the first integral of this system is given by

$$\tau^2 = -e \left[r - 2\mu\sigma(\xi) + \frac{\mu^2 + \rho}{r} \sigma^2(\xi) \right], \quad (2.7)$$

where $\rho = \pm 1$ and $e = \pm 1$.

We consider the following solutions of the system (2.6).

1. **Case I:**

If $r = \mu = 0$ then

$$\tau_1(\xi) = -\frac{1}{e\xi}, \quad \sigma_1(\xi) = \frac{C}{\xi}. \quad (2.8)$$

2. **Case II:**

If $e = 1$, $\rho = -1$ and $r > 0$:

$$\begin{cases} \sigma_1(\xi) = \frac{r \sec(\sqrt{r} \xi)}{1 + \mu \sec(\sqrt{r} \xi)}; & \tau_1(\xi) = \frac{\sqrt{r} \tan(\sqrt{r} \xi)}{1 + \mu \sec(\sqrt{r} \xi)} \\ \sigma_2(\xi) = \frac{r \csc(\sqrt{r} \xi)}{1 + \mu \csc(\sqrt{r} \xi)}; & \tau_2(\xi) = -\frac{\sqrt{r} \cot(\sqrt{r} \xi)}{1 + \mu \csc(\sqrt{r} \xi)}. \end{cases} \quad (2.9)$$

3. **Case III:**

If $e = -1$, $\rho = -1$ and $r > 0$:

$$\begin{cases} \sigma_3 = \frac{r \operatorname{sech}(\sqrt{r} \xi)}{1 + \mu \operatorname{sech}(\sqrt{r} \xi)} \\ \tau_3 = \frac{\sqrt{r} \tanh(\sqrt{r} \xi)}{1 + \mu \operatorname{sech}(\sqrt{r} \xi)}. \end{cases} \quad (2.10)$$

4. **Case IV:**

If $e = -1$, $\rho = 1$ and $r > 0$:

$$\begin{cases} \sigma_4 = \frac{r \operatorname{csch}(\sqrt{r} \xi)}{1 + \mu \operatorname{csch}(\sqrt{r} \xi)} \\ \tau_4 = \frac{\sqrt{r} \coth(\sqrt{r} \xi)}{1 + \mu \operatorname{csch}(\sqrt{r} \xi)}. \end{cases} \quad (2.11)$$

Step 2. Substituting (2.5), along with (2.6) and (2.7) into (2.4) and collecting all terms with the same power in $\sigma^i(\xi)\tau^j(\xi)$, we get a polynomial in the two variables $\sigma(\xi)$ and $\tau(\xi)$. This polynomial has the form

$$\begin{aligned} & a\sigma(\xi)^{m+5} + b\sigma(\xi)^{2m+3} + c\sigma(\xi)^{m+4}\tau(\xi) + d\sigma(\xi)^{3m+1} + \\ & + e\sigma(\xi)^{2m+2}\tau(\xi) + \text{other terms of lower degree} \end{aligned} \quad (2.12)$$

We assume that $m \geq 1$ to avoid trivial solutions. The degrees of the highest terms are $m+5$ (the degree of the terms $a\sigma(\xi)^{m+5}$ and $c\sigma(\xi)^{m+4}\tau(\xi)$), $2m+3$ (the degree of the term $b\sigma(\xi)^{2m+3}$) and $3m+1$ (the degree of the term $d\sigma(\xi)^{3m+1}$). There are two integer values of m for which $3m+2 = 2m+3$ or $3m+1 = m+5$ or $2m+3 = m+5$. These are $m = 1$ and $m = 2$. We are going to find solutions for $m = 1$ (the case $m = 2$ is not considered here). When $m = 1$ solutions have the form $v(\xi) = a_0 + a_1\sigma(\xi) + b_1\tau(\xi)$ and equating in (2.12) the coefficients of every power of $\sigma(\xi)$ and of every term of the form $\sigma^j(\xi)\tau(\xi)$ to zero, we obtain the following algebraic system in the variables $a_0, a_1, b_1, \dots$:

1. $e^7(\mu^2 + \rho)^2 a_1 = 0$
2. $e^8(\mu^2 + \rho)^3 b_1 = 0$
3. $e^5(\mu^2 + \rho)^2(6\mu e^3 - 3\mu e + 4a_1) b_1 = 0$
4. $e^4(\mu^2 + \rho)(2r\mu a_1 e^3 - \mu^2 b_1^2 e - \rho b_1^2 e - r\mu a_1 e + r a_1^2) = 0$
5. $(e-1)(e+1)r(2re^3 - re - 3a_0) b_1^2 = 0$
6. $120r^2 a_1 e^7 - 180r^2 a_1 e^5 - 180r a_0 a_1 e^4 + 61r^2 a_1 e^3 + 150r a_0 a_1 e^2 + 180a_0^2 a_1 e + \lambda a_1 e - 960r\mu b_1^2 e^5 + 1230r\mu b_1^2 e^3 + 720\mu a_0 b_1^2 e^2 - 540r a_1 b_1^2 e^2 - 300r\mu b_1^2 e - 360\mu a_0 b_1^2 + 360r a_1 b_1^2 = 0$
7. $16r^2 \mu a_1 e^7 - 20r^2 \mu a_1 e^5 + 8r^2 a_1^2 e^4 - 12r\mu a_0 a_1 e^4 + 5r^2 \mu a_1 e^3 - 6r^2 a_1^2 e^2 + 6r\mu a_0 a_1 e^2 - 48r\mu^2 b_1^2 e^5 - 16r\rho b_1^2 e^5 + 46r\mu^2 b_1^2 e^3 + 12r\rho b_1^2 e^3 + 12\mu^2 a_0 b_1^2 e^2 - 12r a_0 a_1^2 e - 6r\mu^2 b_1^2 e + 12e^2 \rho a_0 b_1^2 - 36e^2 r\mu a_1 b_1^2 + 12r\mu a_1 b_1^2 = 0$

8. $24r\mu^2a_1e^6 + 8r\rho a_1e^6 - 22r\mu^2a_1e^4 - 6r\rho a_1e^4 - 6\mu^2a_0a_1e^3 - 6\rho a_0a_1e^3 + 3r\mu^2a_1e^2$
 $-32\mu^3b_1^2e^4 - 32\mu\rho b_1^2e^4 + 16r\mu a_1^2e^3 + 17\mu^3b_1^2e^2 + 17\mu\rho b_1^2e^2 - 7r\mu a_1^2e + 6ra_1^3$
 $-18e\mu^2a_1b_1^2 - 18e\rho a_1b_1^2 = 0$
9. $(e-1)(e+1)rb_1(120r^2e^6 - 120r^2e^4 - 180ra_0e^3 + 16r^2e^2 - 180rb_1^2e$
 $+60ra_0e + 180a_0^2 + \lambda) = 0$
10. $b_1(720r^2\mu e^8 - 1320r^2\mu e^6 + 662r^2\mu e^4 - 61r^2\mu e^2 + 2\lambda\mu e^2 - 150r\mu a_0e - \lambda\mu$
 $-720r\mu a_0e^5 + 480r^2a_1e^5 + 840r\mu a_0e^3 - 540r^2a_1e^3 + 360\mu a_0^2e^2 + 90r^2a_1e - 180\mu a_0^2$
 $-720r\mu b_1^2e^3 - 720ra_0a_1e^2 + 540r\mu b_1^2e + 360ra_0a_1) = 0$
11. $b_1(1800r^2\mu^2e^8 - 2880r^2\mu^2e^6 - 480r^2\rho e^6 + 1186r^2\mu^2e^4 + 136r^2\rho e^4 - 75r^2\mu^2e^2 + \lambda\mu^2e^2$
 $+360r^2\rho e^8 - 1080r\mu^2a_0e^5 - 360r\rho a_0e^5 + 960r\mu^2a_0e^3 + 240r\rho a_0e^3 + \lambda\rho e^2 - 90r\mu^2a_0e$
 $+1920r^2\mu a_1e^5 - 1800r^2\mu a_1e^3 + 180\mu^2a_0^2e^2 + 180\rho a_0^2e^2 - 1440r\mu a_0a_1e^2 + 210r^2\mu a_1e$
 $+360r\mu a_0a_1 - 1080r\mu^2b_1^2e^3 - 360r\rho b_1^2e^3 + 540r^2a_1^2e^2 + 540r\mu^2b_1^2e$
 $+180r\rho b_1^2e - 180r^2a_1^2) = 0$
12. $b_1(80r\mu^3e^8 + 48r\mu\rho e^8 - 104r\mu^3e^6 - 56r\mu\rho e^6 + 31r\mu^3e^4 + 13r\mu\rho e^4 - r\mu^3e^2$
 $-24\mu^3a_0e^5 - 24\mu\rho a_0e^5 + 96r\mu^2a_1e^5 + 12\mu^3a_0e^3 + 12\mu\rho a_0e^3 - 66r\mu^2a_1e^3 + 4r\mu^2a_1e$
 $+32r\rho a_1e^5 - 18r\rho a_1e^3 + 36r\mu a_1^2e^2 - 24\mu^2a_0a_1e^2 - 24\rho a_0a_1e^2 + 6\mu^3b_1^2e - 6r\mu a_1^2$
 $-24\mu^3b_1^2e^3 - 24\mu\rho b_1^2e^3 + 6\mu\rho b_1^2e) = 0$
13. $30e^2(\mu^2 + \rho)b_1(60r\mu^2e^6 + 12r\rho e^6 - 56r\mu^2e^4 - 8r\rho e^4 - 6\mu^2a_0e^3 + 9r\mu^2e^2$
 $-6\rho a_0e^3 + 64r\mu a_1e^3 - 6\mu^2b_1^2e - 6\rho b_1^2e - 24r\mu a_1e + 18ra_1^2) = 0$

Step 3. (This is the more difficult step) Solving the previous system for r, μ, a_0, a_1, b_1 with the aid of $\textcircled{R}$ Mathematica 6.0 we get $b_1 = 0$, so solutions have the form $u = a_0 + a_1\sigma(x + \lambda t)$. Tables 1, 2, 3 and 4 show the results of calculations.

a_0	a_1	μ	$u(x, t); \quad \xi = x + \lambda t$
$\frac{1}{60} (5r - \sqrt{5}\sqrt{r^2 - 4\lambda})$	$\frac{1}{2}$	-1	$\frac{1}{60} (5r - \sqrt{5}\sqrt{r^2 - 4\lambda}) + \frac{r \csc(\sqrt{r} \xi)}{2(1 - \csc(\sqrt{r} \xi))}$
$\frac{1}{60} (5r + \sqrt{5}\sqrt{r^2 - 4\lambda})$	$\frac{1}{2}$	-1	$\frac{1}{60} (5r + \sqrt{5}\sqrt{r^2 - 4\lambda}) + \frac{r \csc(\sqrt{r} \xi)}{2(1 - \csc(\sqrt{r} \xi))}$
$\frac{1}{60} (5r + \sqrt{5}\sqrt{r^2 - 4\lambda})$	$-\frac{1}{2}$	1	$\frac{1}{60} (5r + \sqrt{5}\sqrt{r^2 - 4\lambda}) - \frac{r \csc(\sqrt{r} \xi)}{2(1 + \csc(\sqrt{r} \xi))}$
$\frac{\sqrt{\lambda}}{6}$	$-\frac{1}{2}$	1	$\frac{\sqrt{\lambda}}{6} - \frac{\sqrt{\lambda} \csc\left(\sqrt[4]{4\lambda} \xi\right)}{1 + \csc\left(\sqrt[4]{4\lambda} \xi\right)}$

Table 1 : *Solutions for $r > 0$, $e = 1$ and $\rho = 1$.*

a_0	a_1	μ	$u(x, t); \quad \xi = x + \lambda t$
$\frac{1}{60} (-5r - \sqrt{5}\sqrt{r^2 - 4\lambda})$	$-\frac{i}{2}$	$-i$	$\frac{1}{60} (-5r - \sqrt{5}\sqrt{r^2 - 4\lambda}) + \frac{1}{2}r \left(1 + \frac{1}{\csc(\sqrt{-r}\xi) - 1} \right)$
$\frac{1}{60} (-5r + \sqrt{5}\sqrt{r^2 - 4\lambda})$	$-\frac{i}{2}$	$-i$	$\frac{1}{60} (-5r + \sqrt{5}\sqrt{r^2 - 4\lambda}) + \frac{1}{2}r \left(1 + \frac{1}{\csc(\sqrt{-r}\xi) - 1} \right)$
$\frac{1}{60} (-5r + \sqrt{5}\sqrt{r^2 - 4\lambda})$	$\frac{i}{2}$	i	$\frac{1}{60} (-5r + \sqrt{5}\sqrt{r^2 - 4\lambda}) + \frac{1}{2}r \left(1 - \frac{1}{\csc(\sqrt{-r}\xi) + 1} \right)$
$-\frac{5\sqrt{\lambda}}{6} + \frac{\sqrt{\lambda}}{6}$	$\frac{i}{2}$	i	$\frac{\sqrt{\lambda}}{\csc(\sqrt[4]{4\lambda}\xi) + 1}$

Table 2 : *Solutions for $r < 0$, $e = -1$ and $\rho = 1$.*

a_0	a_1	μ	$u(x, t); \quad \xi = x + \lambda t$
$\frac{1}{60} (5r - \sqrt{5}\sqrt{r^2 - 4\lambda})$	$\frac{1}{2}$	-1	$\frac{1}{60} (5r - \sqrt{5}\sqrt{r^2 - 4\lambda}) + \frac{r \sec(\sqrt{r} \xi)}{2 (1 - \sec(\sqrt{r} \xi))}$
$\frac{1}{60} (5r + \sqrt{5}\sqrt{r^2 - 4\lambda})$	$\frac{1}{2}$	-1	$\frac{1}{60} (5r + \sqrt{5}\sqrt{r^2 - 4\lambda}) + \frac{r \sec(\sqrt{r} \xi)}{2 (1 - \sec(\sqrt{r} \xi))}$
$\frac{1}{60} (5r - \sqrt{5}\sqrt{r^2 - 4\lambda})$	$-\frac{1}{2}$	1	$\frac{1}{60} (5r - \sqrt{5}\sqrt{r^2 - 4\lambda}) - \frac{r \sec(\sqrt{r} \xi)}{2 (\sec(\sqrt{r} \xi) + 1)}$
$\frac{1}{60} (5r + \sqrt{5}\sqrt{r^2 - 4\lambda})$	$-\frac{1}{2}$	1	$\frac{1}{60} (5r + \sqrt{5}\sqrt{r^2 - 4\lambda}) - \frac{r \sec(\sqrt{r} \xi)}{2 (\sec(\sqrt{r} \xi) + 1)}$
$-\frac{\sqrt{\lambda}}{6}$	$\frac{1}{2}$	-1	$-\frac{\sqrt{\lambda}}{6} - \frac{1}{2}\sqrt{\lambda} \operatorname{csch}^2\left(\sqrt[4]{\frac{\lambda}{4}} \xi\right)$
$-\frac{\sqrt{\lambda}}{6}$	$-\frac{1}{2}$	1	$-\frac{\sqrt{\lambda}}{6} + \frac{\sqrt{\lambda}}{\cosh\left(\sqrt[4]{4\lambda} \xi\right) + 1}$
$\frac{\sqrt{\lambda}}{6}$	$\frac{1}{2}$	-1	$\frac{\sqrt{\lambda}}{6} + \frac{\sqrt{\lambda} \sec\left(\sqrt[4]{4\lambda} \xi\right)}{1 - \sec\left(\sqrt[4]{4\lambda} \xi\right)}$

Table 3 : *Solutions for $e = 1$ and $\rho = -1$.*

a_0	a_1	μ	$u(x, t); \quad \xi = x + \lambda t$
$\frac{1}{60} (-5r - \sqrt{5}\sqrt{r^2 - 4\lambda})$	$-\frac{1}{2}$	-1	$\frac{1}{60} (-5r - \sqrt{5}\sqrt{r^2 - 4\lambda}) - \frac{r \operatorname{sech}(\sqrt{r} \xi)}{2(1 - \operatorname{sech}(\sqrt{r} \xi))}$
$\frac{1}{60} (-5r + \sqrt{5}\sqrt{r^2 - 4\lambda})$	$-\frac{1}{2}$	-1	$\frac{1}{60} (-5r + \sqrt{5}\sqrt{r^2 - 4\lambda}) - \frac{r \operatorname{sech}(\sqrt{r} \xi)}{2(1 - \operatorname{sech}(\sqrt{r} \xi))}$
$\frac{1}{60} (-5r - \sqrt{5}\sqrt{r^2 - 4\lambda})$	$\frac{1}{2}$	1	$\frac{1}{60} (-5r - \sqrt{5}\sqrt{r^2 - 4\lambda}) + \frac{r \operatorname{sech}(\sqrt{r} \xi)}{2(\operatorname{sech}(\sqrt{r} \xi) + 1)}$
$\frac{1}{60} (-5r + \sqrt{5}\sqrt{r^2 - 4\lambda})$	$\frac{1}{2}$	1	$\frac{1}{60} (-5r + \sqrt{5}\sqrt{r^2 - 4\lambda}) + \frac{r \operatorname{sech}(\sqrt{r} \xi)}{2(\operatorname{sech}(\sqrt{r} \xi) + 1)}$
$-\frac{\sqrt{\lambda}}{6}$	$-\frac{1}{2}$	-1	$-\frac{\sqrt{\lambda}}{6} - \frac{\sqrt{\lambda} \operatorname{sech}(\sqrt[4]{4\lambda} \xi)}{1 - \operatorname{sech}(\sqrt[4]{4\lambda} \xi)}$
$-\frac{\sqrt{\lambda}}{6}$	$\frac{1}{2}$	1	$-\frac{\sqrt{\lambda}}{6} + \frac{\sqrt{\lambda} \operatorname{sech}(\sqrt[4]{4\lambda} \xi)}{\operatorname{sech}(\sqrt[4]{4\lambda} \xi) + 1}$

Table 4 : *Solutions for $r > 0$, $e = -1$ and $\rho = -1$.*

3 Conclusions

In this paper, by using the projective Riccati equation method and the help of symbolic computation system *Mathematica*, we obtain some exact solutions for the equation (1.2). The projective Riccati equation method is more complicated than others methods (for example, the tanh method) in the sense that the algebraic system in many cases demands a lot of time to be solved. On the other hand, this method allows us to obtain some new exact solutions.

References

- [1] R. CONTE & M. MUSETTE , *Link betwen solitary waves and projective Riccati equations*, J. Phys. A Math. **25** (1992), 5609-5623.
- [2] E. INC & M. ERGT, *New Exact Travelling Wave Solutions for Compound KdV-Burgers Equation in Mathematical Physics*, Applied Mathematics E-Notes, **2**(2002), 45-50.
- [3] J. MEI, H. ZHANG & D. JIANG, *New exact solutions for a Reaction-Diffusion equation and a Quasi-Camassa-Holm Equation*, Applied Mathematics E-Notes, **4**(2004), 85-91.
- [4] Z. YAN, *The Riccati equation with variable coefficients expansion algorithm to find more exact solutions of nonlinear differential equation*, MMRC, AMSS, Academis Sinica, Beijing **22**(2003), 275-284.
- [5] Z. YAN, *An improved algebra method and its applications in nonlinear wave euations*, MMRC, AMSS, Academis Sinica, Beijing **22**(2003), 264-274.
- [6] W.HEREMAN, *Symbolic computation of exact solutions expressible in hyperbolic and elliptic functions for nonlinear partial differential and diffential-difference equations*, Journal of Symbolic Computation.
- [7] ALVARO H. SALAS & C. GOMEZ, *El mathematica en la búsqueda de soluciones exactas para ecuaciones diferenciales parciales lineales y no lineales*, Primer Simposio Inter-

nacional del uso de Tecnologías en educación matemática. Universidad Pedagógica Nacional, Bogotá, Colombia. **1**(2005).

- [8] ALVARO H. SALAS, *Some solutions for a type of generalized Sawada-Kotera equation*, Applied Mathematics and Computation, Elsevier Editorial System, Netherlands, in press, available on line from july 17 2007 at <http://www.sciencedirect.com>
- [9] C. GOMEZ, ALVARO H. SALAS, *New exact solutions for the combined sinh-cosh-Gordon equation*, Lecturas matemáticas, volumen especial 2006, Apuntes XV Congreso Nacional de Matemáticas, agosto de 2005, Bogotá, Colombia.
- [10] XUEQIN ZHAO, HONGYAN ZHI, YAXUAN YU AND HONGQING ZHANG, *A new Riccati equation method with symbolic computation to construct new travelling wave solution of nonlinear differential equations*, Applied mathematics and Computation 172 (2006) 24-39.